Patient Quick Start Guide for Aimovig® (erenumab-aooe)

Follow these tips to help you start and stay on track with treatment.

1 Enrolling in copay support

- Eligible, commercially insured patients may **pay as little as $49 per month**.* To enroll in the Aimovig® Copay Card program, visit **aimovig.com/enrollment**

*Eligibility criteria and program maximums apply. Terms and conditions apply. Please see full Terms and Conditions at **aimovig.com/toc**.

Medicare, Medicaid, or no insurance? Visit amgensupportplus.com for potential financial support options

2 Starting on Aimovig®

- Pick up Aimovig® from your local pharmacy or have it delivered to your home. If eligible, ensure your pharmacist has your copay card details
- For supplemental injection support, watch a video on how to self-inject Aimovig®. Always refer to the Instructions for Use
- Choose a day to take Aimovig® each month
- You can request a free sharps disposal container to properly dispose of used autoinjectors by calling 833-AIMOVIG (833-246-6844)

3 Staying on track

- Sign up for Amgen SupportPlus programs for resources to help throughout your treatment experience:

90-Day Start Program
Receive personalized email support specifically designed for your first 3 months of treatment

Migraine Tracker
Track monthly migraine days with a simple daily text and get personal monthly trend reports

Refill Reminders
Receive a text message to remind you when it's time to pick up or refill your Aimovig® prescription

Get started by scanning the QR code or visiting aimovig.com/support-and-resources

The American Headache Society recommends waiting **at least 3 months** before assessing the benefits of anti-CGRP* pathway preventive treatments like Aimovig®.[1]

*CGRP, calcitonin gene-related peptide.

APPROVED USE

Aimovig® (erenumab-aooe) is a prescription medicine used for the preventive treatment of migraine in adults.

IMPORTANT SAFETY INFORMATION

Who should not use Aimovig®?

Do not use Aimovig® if you are allergic to erenumab-aooe or any ingredients in Aimovig®.

Please see additional Important Safety Information on the next page.

aimovig® (erenumab-aooe) injection 70 mg/mL • 140 mg/mL | AMGEN® Support+

IMPORTANT SAFETY INFORMATION (cont'd)

Before starting Aimovig®, tell your healthcare provider (HCP) about all your medical conditions, including if you have high blood pressure, have circulation problems in your fingers and toes, are pregnant or plan to become pregnant, are breastfeeding or plan to breastfeed.

Tell your HCP about all the medicines you take, including any prescription and over-the-counter medicines, vitamins, or herbal supplements.

What are possible side effects of Aimovig®?

Aimovig® may cause serious side effects, including:

- **Allergic reactions.** Allergic reactions, including rash or swelling can happen after receiving Aimovig®. This can happen within hours to days after using Aimovig®. Call your HCP or get emergency medical help right away if you have any of the following symptoms of an allergic reaction: swelling of the face, mouth, tongue or throat, or trouble breathing.
- **Constipation with serious complications.** Severe constipation can happen after receiving Aimovig®. In some cases people have been hospitalized or needed surgery. Contact your HCP if you have severe constipation or constipation associated with symptoms such as severe or constant belly pain, vomiting, swelling of belly or bloating.
- **High blood pressure.** High blood pressure or worsening of high blood pressure can happen after receiving Aimovig®. Contact your healthcare provider if you have an increase in blood pressure.
- **Raynaud's Phenomenon.** A type of circulation problem can worsen or happen after receiving AIMOVIG. Raynaud's phenomenon can lead to your fingers or toes feeling numb, cool, or painful, or changing color from pale, to blue, to red. Contact your healthcare provider if these symptoms occur

The most common side effects of Aimovig® are pain, redness, or swelling at the injection site and constipation.

These are not all of the possible side effects of Aimovig®. Call your HCP for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit **www.fda.gov/medwatch**, or call 1-800-FDA-1088.

Please click here for Aimovig® full Prescribing Information.

Reference: 1. Ailani J, Burch RC, Robbins SR. The American Headache Society Consensus Statement: Update on integrating new migraine treatments into clinical practice. *Headache.* 2021;61:1021-1039.

AMGEN

© 2026 Amgen Inc. All rights reserved. USA-334-86158 11/25

aimovig® (erenumab-aooe) injection 70 mg/mL • 140 mg/mL